

Properties of galaxy disks in hierarchical hydrodynamical simulations: Comparison with observational data

A. Sáiz, R. Domínguez-Tenreiro

*Depto. de Física Teórica, C-XI. Universidad Autónoma de Madrid,
 Madrid, E-28049, Spain*

P. B. Tissera

I.A.F.E., Casilla de Correos 67, Suc. 28, Buenos Aires, 1428, Argentina

S. Courteau

*University of British Columbia, Dept. of Physics and Astronomy,
 Vancouver, BC, Canada V6T 1Z1*

Abstract. We analyze the structural and dynamical properties of disk-like objects formed in fully consistent cosmological simulations with an inefficient star formation algorithm. Comparison with data of similar observable properties of spiral galaxies gives satisfactory agreement.

We present results of a detailed comparison between the parameters characterizing the structural and dynamical properties of a sample of 29 simulated disk-like objects (DLOs) and those measured in observed spiral galaxies. The DLOs have been identified in AP3M-SPH fully consistent hierarchical hydrodynamical simulations, where an *inefficient* Schmidt law-like algorithm to model the stellar formation processes has been implemented.

A surface density bulge-disk decomposition was performed on the DLOs, using a double-exponential profile. The resulting bulge and disk scale lengths, R_b and R_d , and their ratio R_b/R_d , are consistent with available data (Courteau, de Jong, & Broeils 1996; de Jong 1996; Moriondo, Giovanelli, & Haynes 1999). See Fig. 1(a).

Our rotation curve analysis and shape modeling uses a parameterization based on the spatial scale $R_{2.2} = 2.2 R_d$ and the velocity (mass) estimate $V_{2.2} = V_{\text{cir}}(R_{2.2})$. These parameters were compared with those measured for the Courteau–Faber sample of bright Sb–Sc field spirals with long-slit H α rotation curves, and the Broeils (1992) compilation of late-type spirals with extended H I rotation curves. Deep surface photometry is available for both samples. In contrast to findings in other fully consistent hydrodynamical simulations (Navarro & Steinmetz 2000 and references therein), we find DLO $V_{2.2}$ values that are consistent with the observational data. This is a consequence of disk formation with conservation of the specific angular momentum j . See Fig. 1(b).

In conclusion, the comparison between DLOs produced in our simulations and observational data allows us to affirm that DLOs have counterparts in today spiral galaxies. This agreement suggests that the process operating in Fall & Efstathiou (1980) standard model for disk formation (i.e., gas cooling and




Figure 1. The scale lengths R_b and R_d (a), and the $V_{2,2}$ velocities versus R_d (b). Filled symbols correspond to DLOs formed in our simulations. Dots in (a) are from 1D decompositions of surface brightness profiles given by Courteau, de Jong, & Broeils (1996) and Courteau (1997). Dots in (b) are data from the Courteau-Faber ($H\alpha$ spectra) sample. Open circles are data from spirals with HI rotation curves from Casertano & van Gorkom (1991).

collapse with j conservation) is also at work in the *quiescent* phases of DLO formation in these simulations. However, *violent* episodes (i.e., interactions and merger events) also occur and play an important role in DLO assembly (Domínguez-Tenreiro, Tissera, & Sáiz 1998). To provide the right conditions for disk regeneration after the last violent episode of DLO assembly, a compact central stellar bulge is needed; this will ensure the axisymmetric character of the gravitational potential well at scales of some kpc at all times, avoiding excessive j losses in violent events. A second condition is necessary: the availability of gas at low z , to form the disk. The good match of our DLO parameters with observational data suggests that our *inefficient* star formation algorithm meets both requirements. This global agreement with observations also represents an important step towards making numerical approaches more widely used for the study of galaxy formation and evolution in a cosmological framework, i.e., from primordial fluctuations.

References

- Broeils, A. H. 1992, Ph.D. thesis, Univ. Groningen
- Casertano, S., & van Gorkom, J. H. 1991, AJ, 101, 1231
- Courteau, S. 1997, AJ, 114, 2402
- Courteau, S., de Jong, R. S., & Broeils, A.H. 1996, ApJ, 457, L73
- de Jong, R. S. 1996, A&A, 313, 45
- Domínguez-Tenreiro, R., Tissera P. B., & Sáiz, A. 1998, ApJ, 508, L123
- Fall, S. M., & Efstathiou, G. 1980, MNRAS, 193, 189
- Moriondo, G., Giovanelli, R., & Haynes, M. P. 1999, A&A, 346, 415
- Navarro, J. F., & Steinmetz, M. 2000, ApJ, 538, 477